

## Physics 406: Homework 5

1. **Two spin systems in contact:** A result which we will use a lot in this course is Sterling's approximation for the logarithm of a factorial, which says that

$$\ln n! \simeq n \ln n - n,$$

where the approximation becomes better and better as  $n \rightarrow \infty$ . We will see how to prove this result in a later lecture. For the moment we just assume it. (If you want to see the proof, it's given in Appendix A of Kittel and Kroemer.)

Now consider the spin system we looked at, composed of  $N$  "spins" that each point either up or down.

- (a) Write down an expression for the multiplicity  $g(N, s)$  of states that have a given value of the spin excess parameter  $s$ , defined by  $2s = N_{\uparrow} - N_{\downarrow}$ . Hence write down an expression for the logarithm of the multiplicity as a function of  $s$ .
- (b) Apply Sterling's approximation and show that for large  $N$

$$\ln g = N \ln N - \left(\frac{1}{2}N + s\right) \ln\left(\frac{1}{2}N + s\right) - \left(\frac{1}{2}N - s\right) \ln\left(\frac{1}{2}N - s\right). \quad (1)$$

- (c) Noting that  $\ln\left(\frac{1}{2}N + s\right) = \ln\left(\frac{1}{2}N\right) + \ln\left(1 + 2s/N\right)$ , expand to *second order* in  $2s/N$  and hence show that

$$g(N, s) \simeq 2^N e^{-2s^2/N}. \quad (2)$$

This is a Gaussian or normal distribution: the binomial distribution becomes a Gaussian distribution for large  $N$ .

- (d) Now suppose we have two identical systems of  $N$  spins. Using Eq. (2), write down an expression for the total multiplicity of both systems together as a function of the spin excesses  $s_1$  and  $s_2$  of the two systems, assuming for the moment that the two systems are not in contact with one another. (To make the calculations easier, you can assume  $N$  is even.) Eliminate  $s_2$  in favor of the total spin excess  $s = s_1 + s_2$  and, by completing the square, show that for a given value of  $s$  the distribution of possible values of  $s_1$  is Gaussian.
- (e) Now suppose that both systems are in a magnetic field of intensity  $B$ , so that they have energies  $U_1 = -2mBs_1$  and  $U_2 = -2mBs_2$ , where  $m$  is the dipole moment of each spin. If the two systems are now in contact with one another, so that energy can flow between them, then  $s_1$  and  $s_2$  are no longer fixed, but the total energy must be conserved, so  $s = s_1 + s_2$  is constant. What is the most likely value of  $s_1$ ?

2. **Entropy of a set of harmonic oscillators:** Consider the ensemble that we discussed earlier of  $N$  quantum harmonic oscillators. Each can have energy  $\epsilon_s = s\hbar\omega$ , where  $\hbar$  and  $\omega$  are constants and  $s$  is a non-negative integer. As we showed in class, the multiplicity  $g(N, n)$  of states of the entire ensemble that have total internal energy  $U = n\hbar\omega$  is given by

$$g(N, n) = \binom{N-1+n}{N-1}.$$

- (a) Write  $g(N, n)$  in a form involving factorials and hence write down the dimensionless entropy  $\sigma$  of the system.

- (b) When  $N$  is large we can, to a good approximation, replace  $N - 1$  by  $N$ . Apply Sterling's approximation and derive an approximate expression for  $\sigma$  for large  $N$ .
- (c) Recalling the definition of the temperature  $\tau$  in energy units,  $\tau = \partial U / \partial \sigma$ , differentiate to get an expression for  $\tau$  in terms of the internal energy. (You have to consider  $n$  to be a continuous variable to do this calculation, which is strictly speaking not correct—it is an integer. Later in the course we'll see a better derivation of this result that doesn't require us to do this kludge.)
- (d) Rearrange to show that

$$U = \frac{N\hbar\omega}{\exp(\hbar\omega/\tau) - 1}.$$

This is the internal energy of a set of  $N$  harmonic oscillators and, as we will show, is also the correct expression for the energy of a set of bosons (e.g., photons) in a quantum gas. From this expression we will later derive the famous black-body radiation spectrum of Rayleigh and Planck.

- (e) What is the heat capacity of the system?

3. **Partition function of a simple system:** Suppose a simple system has states with three energies,  $-\varepsilon$ ,  $0$ , and  $+\varepsilon$ . The multiplicities of the states are  $g(-\varepsilon) = 1$ ,  $g(0) = 2$ , and  $g(\varepsilon) = 1$ . The system is put in contact with a thermal reservoir at temperature  $\tau$  (in energy units) and allowed to come to equilibrium.

- (a) Calculate the partition function  $Z$  of the system.
- (b) Calculate the average internal energy of the system as a function of temperature.
- (c) Show that the heat capacity is

$$C = \frac{\varepsilon^2}{\tau^2 [1 + \cosh(\varepsilon/\tau)]}.$$